


Provided for non-commercial research and education use.
Not for reproduction, distribution or commercial use.



This article appeared in a journal published by Elsevier. The attached copy is furnished to the author for internal non-commercial research and education use, including for instruction at the authors institution and sharing with colleagues.

Other uses, including reproduction and distribution, or selling or licensing copies, or posting to personal, institutional or third party websites are prohibited.

In most cases authors are permitted to post their version of the article (e.g. in Word or Tex form) to their personal website or institutional repository. Authors requiring further information regarding Elsevier's archiving and manuscript policies are encouraged to visit:

<http://www.elsevier.com/copyright>



Contents lists available at ScienceDirect

Computers & Geosciences

journal homepage: www.elsevier.com/locate/cageo

Short note

Release of a 10-m-resolution DEM for the Italian territory: Comparison with global-coverage DEMs and anaglyph-mode exploration via the web

Simone Tarquini^{a,*}, Stefano Vinci^b, Massimiliano Favalli^a, Fawzi Doumaz^b,
Alessandro Fornaciai^a, Luca Nannipieri^a^a Istituto Nazionale di Geofisica e Vulcanologia, Sezione di Pisa, Via della Faggiola 32, 56126 Pisa, Italy^b Istituto Nazionale di Geofisica e Vulcanologia, Centro Nazionale Terremoti, Via di Vigna Murata 605, 00143 Roma, Italy

ARTICLE INFO

Article history:

Received 22 December 2010

Received in revised form

25 March 2011

Accepted 30 April 2011

Available online 18 May 2011

Keywords:

DEM

webGIS

Anaglyph

Italy

ABSTRACT

The 10-m-resolution TINITALY/01 DEM (Tarquini et al., 2007) is compared with the two, coarser-resolution, global-coverage, spaceborne-based SRTM and ASTER DEMs and with a high-resolution, LIDAR-derived DEM. Afterwards, we presented a webGIS which allows to explore a 10-m-resolution anaglyph layer showing the landforms of the whole Italian territory in 3D. The webGIS (<http://tinality.pi.ingv.it/>) is open to the public, and can be used to carry out a preliminary analysis of landforms. The TINITALY/01 DEM is available for scientific purposes on the basis of a research agreement (see the above website or write to tinality@pi.ingv.it).

© 2011 Elsevier Ltd. All rights reserved.

1. Introduction

Digital elevation models (DEMs) are fundamental in environmental and morphological studies. DEMs are obtained from a variety of sources and generated in several ways. Nowadays, several elevation datasets are available for free (e.g., Shuttle Radar Topographic Mission (SRTM), <http://www.jpl.nasa.gov/srtm>; ASTER, <http://asterweb.jpl.nasa.gov/>), and the choice of the DEM which better suits the target of the study is crucial (e.g., Hubbard et al., 2007). TINITALY/01 is currently the most accurate DEM covering the whole Italian territory (Tarquini et al., 2007). This DEM was created by using heterogeneous elevation datasets obtained from existing digital cartography.

Human vision provides a 3D perception of the world. Computers can recreate such a capability in a 2D monitor by using several techniques. Anaglyphs are a simple but effective method to convey the depth perception which allows a 3D vision of a DEM. A free webGIS to explore an anaglyph representation of TINITALY/01 is presented, and the whole DEM is now downloadable for scientific purposes.

2. The TINITALY/01 DEM

TINITALY/01 is a DEM in triangular irregular network format (TIN) created for the whole Italian territory in the UTM 32 WGS 84




Fig. 1. Shaded relief image with color table of the TINITALY/01 DEM with the tiles composing the elevation dataset. For the European countries we used the SRTM DEM, while the bathymetry is from the GEBCO dataset (<http://www.gebco.net/>). Ticks mark UTM 32 WGS 84-km coordinates (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article).

* Corresponding author. Tel.: +39 050 8311932; fax: +39 050 8311942.
E-mail address: tarquini@pi.ingv.it (S. Tarquini).

coordinate system (Tarquini et al., 2007). The DEM was obtained from heterogeneous vector datasets, mostly consisting in elevation contour lines and elevation points from several sources. The input vector database was carefully cleaned up to obtain a seamless TIN derived by using the DEST algorithm (Favalli and Pareschi, 2004).

The whole TINITALY/01 DEM was converted in grid format (10-m cell size) according to a tiled structure composed of 193, 50-km side square elements (Fig. 1). The grid database consists of more than 3 billions of cells and occupies ~ 12 GB of disk memory.

3. The TINITALY/01 DEM vs global-coverage DEMs

In recent years, an increasing number of elevation datasets covering entire countries are available to the public. Examples are

the 10- or 30-m resolution National Elevation Dataset for the United States (NED; <http://seamless.usgs.gov>), the 10-m resolution DEM for Japan (Geographical Survey Institute of Japan; <http://fgd.gsi.go.jp/>), the 5- or 10-m resolution DEM for the UK (<http://www.neodc.rl.ac.uk/>), or else the 50-m resolution BD ALTI DEM for France. Here we briefly compare the TINITALY/01 DEM with the two spaceborne-based DEMs with an almost global coverage recently released via the web: the 90-m resolution SRTM and the 30-m resolution ASTER elevation datasets. These two DEMs cover also the whole Italian territory.

At the beginning of this century a quasi-global elevation dataset from the SRTM was released (Rabus et al., 2003). It consists of a 90-m cell size DEM which has been shown to be appropriate for morphometric analyses over large areas or to focus on the broad morphology of features.




Fig. 2. Comparison of the elevation profiles obtained by using a LIDAR-derived topography, the TINITALY/01 grid, the ASTER, and the SRTM elevation datasets.

The ASTER DEM with a 30-m cell size has been released more recently (Hayakawa et al., 2008). The ASTER has a higher resolution with respect to the SRTM dataset. Nevertheless, this fact does not necessarily imply a higher DEM quality, because, as an example, the ASTER DEM is affected by the limits of the ASTER sensor, which is locally disturbed by clouds (Hubbard et al., 2007).

Tarquini et al. (2007) carried out a comprehensive assessment of the accuracy of the TINITALY/01 DEM, finding a root mean square error in elevation (RMSE_z) between 0.8 and 6.0 m. Rabus et al. (2003), for the SRTM elevation dataset, found an absolute vertical accuracy of ± 16 m at 90% confidence, while Hirano et al. (2003) found an RMSE_z for the ASTER DEM within the design specification of 7–50 m. For both ASTER and SRTM DEMs, the error increases as the topography become rougher. As an example, for the ASTER DEM, Kaab (2002) found an RMSE_z of 60 m in rough high mountains and 18 m in moderately mountainous terrains.

An in-depth comparison is beyond the scope of the present work; nevertheless, we selected an example to highlight the characteristics of the TINITALY/01 DEM with respect to the two global-coverage SRTM and ASTER DEMs. We selected Mount Etna (Sicily, Italy), where a very high-resolution, LIDAR-derived topography was available for comparison (Favalli et al., 2009). In this area, the TINITALY/01 DEM showed an RMSE_z of 1.98 m (Neri et al., 2008), while we calculated an RMSE_z of 7.78 and 9.36 m for the SRTM and ASTER DEMs, respectively. Fig. 2 shows the elevation profiles obtained from the LIDAR-derived DEM, the 10-m cell size TINITALY/01 grid, and ASTER and SRTM DEMs. The two spaceborne-based DEMs are too coarse to precisely reveal fundamental features such as cone craters and cone outlines, while the TINITALY/01 DEM provides the necessary detail.

4. Anaglyph-mode navigation of the DEM and related webGIS

The human brain is equipped to produce a 3D perception of an observed scene by coupling two images acquired by the eyes from a slightly different point of view. This is the basic principle of the stereo vision, widely applied for the examination of couples of aerial stereoscopic photographs. The observation in stereo mode conveys a deeper understanding of landscape morphologies, and can fruitfully support the detection of subtle features. Add-on modules enabling 3D vision in anaglyph mode have been recently made available for both freeware and commercial programs (e.g., Google Earth and ESRI ArcGIS), suggesting the validity of this method. Here we use a custom software (Tarquini and Favalli, 2011) which produces full-resolution anaglyph images from arbitrarily large DEMs, to obtain a seamless anaglyph layer of the entire TINITALY/01 database at 10-m resolution.

The ESRI ArcGIS server is used to set up a webGIS (<http://tinality.pi.ingv.it/>), where the described 10-m resolution anaglyph layer is freely accessible for navigation (Fig. 3). The viewpoints for anaglyph creation are set symmetrically to a nadiral axis pointing toward the center of Italy, at an almost infinite distance, obtaining a very low distortion. The anaglyph layer is geo-referenced in the UTM WGS 84 zone 32 projection system. To view the anaglyphs, users must wear anaglyph glasses with a green (or blue) left lens and a red right lens, but we set up also a layer of shaded relief images with an elevation color ramp.

5. Concluding remarks

The 10-m resolution TINITALY/01 DEM (Tarquini et al., 2007) is compared with the two, coarser-resolution, global-coverage,




Fig. 3. Example of anaglyph image of the Majella Mountain (Abruzzo region, Italy), obtained from the TINITALY/01 DEM. Ticks mark UTM 32 WGS 84-km coordinates (For interpretation of the references to color in this figure (color mentioned in the text), the reader is referred to the web version of this article).

spaceborne-based SRTM and ASTER DEMs and with a high resolution LIDAR-derived DEM. Afterward, we presented a web-GIS which allows the exploration of a 10-m resolution anaglyph layer showing the landforms of the whole Italian territory in 3D. The webGIS (<http://tinality.pi.ingv.it/>) is open to the public, and can be used to carry out a preliminary analysis of landforms. The TINITALY/01 DEM is now available for scientific purposes on the basis of a research agreement (see the above website or write to tinality@pi.ingv.it).

References

- Favalli, M., Fornaciai, A., Pareschi, M.T., 2009. LIDAR strip adjustment: application to volcanic areas. *Geomorphology* 111, 123–135.
- Favalli, M., Pareschi, M.T., 2004. Digital elevation model construction from structured topographic data: the DEST algorithm. *Journal of Geophysical Research* 109, F04004.
- Hayakawa, Y.S., Oguchi, T., Lin, Z., 2008. Comparison of new and existing global digital elevation models: ASTER G-DEM and SRTM-3. *Geophysical Research Letters* 35, L17404.
- Hirano, A., Welch, R., Lang, W., 2003. Mapping from ASTER stereo image data: DEM validation and accuracy assessment. *ISPRS Journal of Photogrammetry and Remote Sensing* 57, 356–370.
- Hubbard, B.E., Sheridan, M.F., Carrasco-Núñez, G., Díaz-Castellón, R., Rodríguez, S.R., 2007. Comparative lahar hazard mapping at Volcan Citlaltépetl, Mexico using SRTM, ASTER and DTED-1 digital topographic data. *Journal of Volcanology and Geothermal Research* 160, 99–124.
- Kaab, A., 2002. Combination of SRTM3 and repeat ASTER data for deriving alpine glacier flow velocities in the Bhutan Himalaya. *Remote Sensing of Environment* 94, 463–474.
- Neri, M., Mazzarini, F., Tarquini, S., Bisson, M., Isola, I., Behncke, B., Pareschi, M.T., 2008. The changing face of Mount Etna's summit area documented with Lidar technology. *Geophysical Research Letters* 35, L09305.
- Rabus, B., Eineder, M., Roth, A., Bamler, R., 2003. The shuttle radar topography mission—a new class of digital elevation models acquired by spaceborne radar. *ISPRS Journal of Photogrammetry and Remote Sensing* 57, 241–262.
- Tarquini, S., Favalli, M., 2011. Capturing full resolution perspective and stereo views of large DEMs. *Rapporto Tecnico INGV*, 181.
- Tarquini, S., Isola, I., Favalli, M., Mazzarini, F., Bisson, M., Pareschi, M.T., Boschi, E., 2007. TINITALY/01: a new triangular irregular network of Italy. *Annals of Geophysics* 50, 407–425.